Humeral Bone Loss in Revision Total Shoulder Arthroplasty: the Proximal Humeral Arthroplasty Revision Osseous inSufficiency (PHAROS) Classification System

Peter N, Chalmers MD, Anthony A. Romeo MD, Gregory P. Nicholson MD, Pascal Boileau MD, PhD, Jay D. Keener MD, James M. Gregory MD, Dane H. Salazar MD, Robert Tashjian MD

Clin Orthop Relat Res. 2019 Feb; 477(2): 432-441.

The abstract can be found at: doi: 10.1097/CORR.000000000000590

When performing revision shoulder arthroplasty, it is common to encounter humeral bone loss. Anticipating humeral defects prior to performing revision shoulder arthroplasty allows the revising surgeon to plan for fixation and stability of the revision humeral prosthesis. Until this publication, there was no known validated classification system for humeral bone loss in revision shoulder arthroplasty. Creation and utilization of a classification system for humeral arthroplasty revision could assist the surgeon in diagnosing, treating and with a final prognosis. The PHAROS system was devised to help surgeons classify humeral defects and to build a strategy for revision humeral arthroplasty where considerations like the need for allograft, an appropriate revision stem length, potential need for proximal humeral replacement in addition to likelihood of intraoperative humeral related complications such as greater tuberosity fractures.

This retrospective, multi-part study, included high-volume, experienced surgeons to review radiographic and clinical histories of patients with revision shoulder arthroplasty from November 2006 to January 2018.

KEY TAKEAWAYS

- Radiographic images and clinical histories were necessary for evaluation. Pre-revision images included AP, Grashey, and lateralview radiographs in addition to patient's clinical history.
- Upon review, revised patients were excluded for the following criteria: no humeral component revision, no new humeral placement, no previous humeral component and where a platform system was utilized but not revised.
- A total of 106 patients with 108 shoulders for revision were included in the cohort.
- Three types of bone loss were classified: Type 1, Type 2 and Type 3.
- Type 1 bone loss includes epiphyseal bone loss with epiphysis including calcar, tuberosities and the articular surface. Type 2 bone loss includes metaphyseal bone loss with the metadiaphysis above the insertion of the deltoid and Type 3 includes diaphyseal bone loss below the deltoid insertion point.
- The PHAROS classification also includes evaluation of cortical walls, condition of the calcar and compromise of the Greater Tuberosity.
- In Type 1 and Type 2 bone loss, use of greater tuberosity fixation for intraoperative fractures was commonly utilized. In Type 2 and Type 3 bone loss, structural humeral bone grafting was commonly utilized and in Type 3, where diaphyseal bone loss was greatest, the most treatment was proximal humeral replacement or total humeral replacement.
- In conclusion, the PHAROS classification system was shown to be helpful in planning operative treatments, but this study does not include data on clinical outcomes, revision rates or rates of humeral loosening.

Proper surgical procedures and techniques are the responsibility of the medical professional. This material is furnished for information purposes only. Each surgeon must evaluate the appropriateness of the material based on his or her personal medical training and experience. Prior to use of the implant system, the surgeon should refer to the product package insert for complete warnings, precautions, indications, contraindications, and adverse effects. Package inserts are also available by contacting Wright.Contact information can be found in this document and the package insert.